



# CIE A Level Chemistry



Your notes

## 22.2 Mass Spectrometry

### Contents

- \* Mass Spectrometry
- \* Isotopic Abundance & Relative Atomic Mass
- \* Molecular Ion Peak & Fragmentation
- \* The M+1 & M+2 Peaks




Your notes

## Mass Spectrometry


### Interpreting Mass Spectra

- Mass spectroscopy is an analytical technique used to identify unknown compounds
- The molecules in the small sample are **bombarded** with high energy electrons which can cause the molecule to lose an electron
- This results in the formation of a positively charged **molecular ion** with one unpaired electron
  - One of the electrons in the pair has been **removed** by the beam of electrons



- $\text{MOLECULE}^+\bullet$  represents the molecular ion
- The molecular ion can further **fragment** to form new ions, molecules, and radicals

#### Fragmentation of a molecule in mass spectroscopy



Copyright © Save My Exams. All Rights Reserved

*The same molecule can produce several different fragments in mass spectroscopy*

- These **fragmentation** ions are **accelerated** by an electric field




Your notes

- Based on their mass (**m**) to charge (**e**) ratio, the fragments of ions are then separated by deflecting them into the **detector**
  - For example, an ion with mass 16 and charge 2+ will have a  $m/e$  value of 8
- The smaller and more positively charged fragment ions will be **detected** first as they will get **deflected** the most and are more attracted to the **negative pole** of the magnet
- Each fragment corresponds to a specific **peak** with a particular  $m/e$  value in the mass spectrum
- The **base peak** is the peak corresponding to the most **abundant** ion

## Isotopes

- Isotopes are different atoms of the **same element** that contain the same number of **protons** and **electrons** but a different number of **neutrons**.
  - These are atoms of the same **elements** but with different mass number
  - For example, Cl-35 and Cl-37 are isotopes as they are both atoms of the same element (chlorine, Cl) but have a different mass number (35 and 37 respectively)
- Mass spectroscopy can be used to find the **relative abundance** of the isotopes experimentally
- The **relative abundance** of an isotope is the proportion of one particular isotope in a mixture of isotopes found in nature
  - For example, the relative abundance of Cl-35 and Cl-37 is 75% and 25% respectively
  - This means that in nature, 75% of the chlorine atoms is the Cl-35 isotope and 25% is the Cl-37 isotope
- The **heights** of the peaks in mass spectroscopy show the proportion of each isotope present

Example mass spectrum of boron



*The peak heights show the relative abundance of the boron isotopes: boron-10 has a relative abundance of 19.9% and boron-11 has a relative abundance of 80.1%*



Your notes

 **Worked example**

In a sample of iron, the ions  $^{54}\text{Fe}^{2+}$  and  $^{56}\text{Fe}^{3+}$  are detected.

Calculate the  $m/e$  value ratio and determine which ion is deflected more inside the spectrometer.

**Answer:**

- $m/e (^{54}\text{Fe}^{2+}) = \frac{54}{2} = 27$
- $m/e (^{56}\text{Fe}^{3+}) = \frac{56}{3} = 18.7$   
 $= 19$

 **Examiner Tip**

A small  $m/e$  value corresponds to fragments that are either **small** or have a **high positive charge** or a combination of **both**



Your notes

## Isotopic Abundance & Relative Atomic Mass

### Calculating Relative Atomic Mass

- Isotopes are different atoms of the **same element** that contain the same number of **protons** and electrons but a different number of **neutrons**.
  - These are atoms of the same **elements** but with different mass numbers
- Because of this, the mass of an element is given as **relative atomic mass ( $A_r$ )** by using the average mass of the isotopes
- The relative atomic mass of an element can be calculated by using the **relative abundance** values
  - The relative abundance of an isotope is either given or can be read off the mass spectrum
- $$A_r = \frac{(\text{relative abundance}_{\text{isotope 1}} \times \text{mass}_{\text{isotope 1}}) + (\text{relative abundance}_{\text{isotope 2}} \times \text{mass}_{\text{isotope 2}}) \text{ etc.}}{100}$$

#### Worked example

Calculate the relative atomic mass,  $A_r$ , of oxygen to 2 d.p.

Isotope	Percentage abundance
$^{16}\text{O}$	99.76
$^{17}\text{O}$	0.04
$^{18}\text{O}$	0.20

Answer:


- $$A_r = \frac{(99.76 \times 16) + (0.04 \times 17) + (0.20 \times 18)}{100}$$
- $A_r = 16.0044$
- $A_r = \mathbf{16.00}$  (to 2 d.p.)



Your notes

### Worked example

Calculate the relative atomic mass of boron using its mass spectrum, to 2 d.p.



Answer:

- $A_r = \frac{(19.9 \times 10) + (80.1 \times 11)}{100}$
- $A_r = 10.801$
- $A_r = \mathbf{10.80}$  (to 2 d.p)



Your notes

## Molecular Ion Peak & Fragmentation

### Mass Spectrometry: Deducing Molecular Formula

- Each **peak** in the mass spectrum corresponds to a certain **fragment** with a particular  **$m/e$**  value
- The peak with the highest  **$m/e$**  value is the molecular ion ( **$M^+$** ) peak which gives information about the **molecular mass** of the compound
- The molecular ion is the entire molecule that has lost one electron when bombarded with a beam of electrons




- The  **$[M+1]$**  peak is a smaller peak which is due to the natural abundance of the isotope carbon-13
- The height of the  **$[M+1]$**  peak for a particular ion depends on how many carbon atoms are present in that molecule; the more carbon atoms, the larger the  **$[M+1]$**  peak is
  - For example, the height of the  **$[M+1]$**  peak for an hexane (containing six carbon atoms) ion will be greater than the height of the  **$[M+1]$**  peak of an ethane (containing two carbon atoms) ion



Your notes

### Worked example

Determine whether the following mass spectrum corresponds to propanal or butanal.



Copyright © Save My Exams. All Rights Reserved

#### Answer:

- The mass spectrum corresponds to propanal as the molecular ion peak is at  $m/e = 58$
- Propanal arises from the  $\text{CH}_3\text{CH}_2\text{CHO}^+$  ion which has a molecular mass of 58
- Butanal arises from the  $\text{CH}_3\text{CH}_2\text{CH}_2\text{CHO}^+$  ion which has a molecular mass of 72



Your notes

## Identifying Molecules using Fragmentation

- The molecular ion peak can be used to identify the **molecular mass** of a compound
- However, different compounds may have the same molecular mass
- To further determine the structure of the unknown compound, **fragmentation** is used
- Fragments may appear due to the formation of **characteristic fragments** or the **loss of small molecules**
  - For example, a peak at 29 is due to the characteristic fragment  $\text{C}_2\text{H}_5^+$
  - Loss of small molecules gives rise to peaks at 18 ( $\text{H}_2\text{O}$ ), 28 ( $\text{CO}$ ), and 44 ( $\text{CO}_2$ )

## Alkanes

- Simple alkanes are fragmented in mass spectroscopy by breaking the C-C bonds
- $m/e$**  values of some of the common alkane fragments are given in the table below


**$m/e$  values of fragments table**

Fragment	$m/e$
$\text{CH}_3^+$	15
$\text{C}_2\text{H}_5^+$	29
$\text{C}_3\text{H}_7^+$	43
$\text{C}_4\text{H}_9^+$	57
$\text{C}_5\text{H}_{11}^+$	71
$\text{C}_6\text{H}_{13}^+$	85

**Mass spectrum showing fragmentation of alkanes**



Your notes



*Straight chain alkanes show characteristic peaks*


## Halogenoalkanes

- Halogenoalkanes often have multiple peaks around the molecular ion peak
- This is caused by the fact that there are different isotopes of the halogens

### Mass spectrum for a bromine containing species



Your notes



**Mass spectra show different halogen isotopes in the molecular ion**



## Alcohols

- Alcohols often tend to lose a **water molecule** giving rise to a peak at **18 below the molecular ion**
- Another common peak is found at  $m/e$  value 31 which corresponds to the  $\text{CH}_2\text{OH}^+$  fragment
- For example, the mass spectrum of propan-1-ol shows that the compound has fragmented in four different ways:
  - Loss of  $\text{H}^\bullet$  to form a  $\text{C}_3\text{H}_7\text{O}^+$  fragment with  $m/e = 59$
  - Loss of a water molecule to form a  $\text{C}_3\text{H}_6^+$  fragment with  $m/e = 42$
  - Loss of a  $\cdot\text{C}_2\text{H}_5$  to form a  $\text{CH}_2\text{OH}^+$  fragment with  $m/e = 31$
  - And the loss of  $\cdot\text{CH}_2\text{OH}$  to form a  $\text{C}_2\text{H}_5^+$  fragment with  $m/e = 29$

### Mass spectrum of propan-1-ol



Your notes



Copyright © Save My Exams. All Rights Reserved

**Mass spectrum showing the fragmentation patterns in propan-1-ol (alcohol)**



Your notes


### Worked example

Which of the following statements about the mass spectrum of  $\text{CH}_3\text{Br}$  is correct?

- A. There is one peak for the molecular ion with an  $m/e$  value of 44
- B. There is one peak for the molecular ion with an  $m/e$  value of 95
- C. The last two peaks have abundances in the ratio 3:1 and occur at  $m/e$  values of 94 and 96
- D. The last two peaks are of equal size and occur at  $m/e$  values of 94 and 96

**Answer:**

- The correct answer is **D** as bromomethane ( $\text{CH}_3\text{Br}$ ) will fragment into 3 peaks
  - $\text{CH}_3^{81}\text{Br} \rightarrow [\text{CH}_3^{81}\text{Br}]^+ + e^-$  at  $m/e$  96
  - $\text{CH}_3^{79}\text{Br} \rightarrow [\text{CH}_3^{79}\text{Br}]^+ + e^-$  at  $m/e$  94
  - $\text{CH}_3\text{Br} \rightarrow [\text{CH}_3]^+ + \cdot\text{Br}$  at  $m/e$  15
- The last two peaks (which correspond to the molecular ion peak) therefore are equal in size and occur at  $m/e$  values of 94 and 96





Your notes

 **Worked example**

Which alcohol is not likely to have a fragment ion at  $m/e$  at 43 in its mass spectrum?

- A.  $(\text{CH}_3)_2\text{CHCH}_2\text{OH}$
- B.  $\text{CH}_3\text{CH}(\text{OH})\text{CH}_2\text{CH}_2\text{CH}_3$
- C.  $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2\text{OH}$
- D.  $\text{CH}_3\text{CH}_2\text{CH}(\text{OH})\text{CH}_3$

**Answer:**

- The correct answer is **D** because a line at  $m/e = 43$  corresponds to an ion with a mass of 43 for example:
  - $[\text{CH}_3\text{CH}_2\text{CH}_2]^+$
  - $[(\text{CH}_3)_2\text{CH}]^+$
- 2-butanol is not likely to have a fragment at  $m/e = 43$  as it does not have either of these fragments in its structure



Your notes

## The M+1 & M+2 Peaks

### Determine Number of Carbon Atoms Using M+1 Peak

- The **[M+1]** peak is caused by the presence of the carbon-13 ( $^{13}\text{C}$ ) isotope in the molecule
- Carbon-13 makes up approximately 1.1% of all carbon atoms
- Therefore, the [M+1] peak is much smaller than the M peak as the isotope is less common
  - The ratio of  $^{13}\text{C}$  to  $^{12}\text{C}$  is approximately 1:99
- Thus, the greater the number of carbon atoms present in a molecule the greater the height of the [M+1] peak
- The number of carbon atoms, **n**, in a compound can be deduced using the [M+1] peak and the following formula:


$$n = \frac{100 \times \text{abundance of } [M+1]}{1.1 \times \text{abundance of } M^+ \text{ ion}}$$



Your notes

### Worked example

Determine the number of carbon atoms of compound **X** with the following mass spectrum:



#### Answer:

- The  $M^+$  ion peak is at  $m/e$  58 with a relative abundance of around 85
- The  $[M+1]$  peak is at  $m/e$  59 with a relative abundance of 3
- Therefore, the number of carbon atoms ( $n$ ) is:

$$n = \frac{100 \times 3}{1.1 \times 85} = 3.21$$

- There are therefore 3 carbon atoms present in compound **X**



Your notes

## Detecting Bromine & Chlorine Atoms Using M+2 Peak

- The presence of bromine or chlorine atoms in a compound gives rise to a **[M+2]** and possibly **[M+4]** peak



### Chlorine

- Chlorine exists as two isotopes,  $^{35}\text{Cl}$  and  $^{37}\text{Cl}$
- A compound containing **one** chlorine atom will therefore have two molecular ion peaks due to the two different isotopes it can contain
  - $^{35}\text{Cl} = \mathbf{M^+}$  peak
  - $^{37}\text{Cl} = \mathbf{[M+2]}$  peak
  - The ratio of the peak heights is 3:1 (as the relative abundance of  $^{35}\text{Cl}$  is 3x greater than that of  $^{37}\text{Cl}$ )
- A compound containing **two** chlorine atoms will have three molecular ion peaks due to the different combinations of chlorine isotopes they can contain
  - $^{35}\text{Cl} + ^{35}\text{Cl} = \mathbf{M^+}$  peak
  - $^{35}\text{Cl} + ^{37}\text{Cl} = \mathbf{[M+2]}$  peak
  - $^{37}\text{Cl} + ^{37}\text{Cl} = \mathbf{[M+4]}$  peak
  - The ratio of the peak heights is **9:6:1**

### Mass spectra of chlorine containing compounds



Your notes



Mass spectrum of compounds containing one chlorine atom (1) and two chlorine atoms (2)

## Bromine



- Bromine too exists as two isotopes,  $^{79}\text{Br}$  and  $^{81}\text{Br}$
- A compound containing **one** bromine atom will have two molecular ion peaks
  - $^{79}\text{Br} = M^+$  peak
  - $^{81}\text{Br} = [M+2]$  peak
  - The ratio of the peak heights is 1:1 (they are of **similar** heights as their relative abundance is the same!)
- A compound containing **two** bromine atoms will have three molecular ion peaks

- $^{79}\text{Br} + ^{79}\text{Br} = \text{M}^+$  peak
- $^{79}\text{Br} + ^{81}\text{Br} = [\text{M}+2]$  peak
- $^{81}\text{Br} + ^{81}\text{Br} = [\text{M}+4]$  peak
- The ratio of the peak heights is 1:2:1



Your notes

### Mass spectra of bromine containing compounds



Copyright © Save My Exams. All Rights Reserved

Mass spectrum of compounds containing one bromine atom (1) and two bromine atoms (2)




Your notes

### Worked example

Two stable isotopes of bromine have relative masses of 79 and 81

Which is the correct pattern of peaks in the mass spectrum of molecular bromine?



**Answer:**

- The correct answer is **D**
- Bromine is a diatomic molecule there will be 5 peaks on the mass spectrum of bromine
- Bromine consists of molecules, not individual atoms
- When bromine is passed through the mass spectrometer, an electron is given off to give the **molecular ion,  $\text{Br}_2^+$**
- Some of these will fragment to make  $\text{Br} + \text{Br}^+$ 
  - $\text{Br}_2^+ \rightarrow \text{Br} + \text{Br}^+$
- The  $\text{Br}$  atom passes through the machine, and the  $\text{Br}^+$  ions will give lines at 79 and 81
- There will also be a line for the unfragmented  $\text{Br}_2^+$  ion
- This will give 3 molecular ion peaks
  - $\text{Br}_2^+$  ion containing the isotopes  $79 + 79 = 158$
  - $\text{Br}_2^+$  containing the isotopes  $79 + 81 = 160$
  - $\text{Br}_2^+$  containing the isotopes  $81 + 81 = 162$