

AS Level Chemistry A

H032/02 Depth in Chemistry

Friday 9 June 2017 – Afternoon

Time allowed: 1 hour 30 minutes

You must have:

 the Data Sheet for Chemistry A (sent with general stationery)

You may use:

· a scientific or graphical calculator



First name				
Last name				
Centre number		Candid		

INSTRUCTIONS

- Use black ink. HB pencil may be used for graphs and diagrams.
- Complete the boxes above with your name, centre number and candidate number.
- Answer all the questions.
- Write your answer to each question in the space provided. If additional answer space
 is required, you should use the lined page(s) at the end of the booklet. The question
 number(s) must be clearly shown.
- Do not write in the barcodes.

INFORMATION

- The total mark for this paper is 70.
- The marks for each question are shown in brackets [].
- Quality of extended responses will be assessed in questions marked with an asterisk (*).
- This document consists of 16 pages.



Answer **all** the questions.

1	Bar	ium d	combines with oxygen, chlorine and nitrogen to form ionic compounds.
	(a)	Bar	ium oxide, BaO, has a giant ionic lattice structure.
		(i)	State what is meant by the term ionic bond.
			[1]
		(ii)	Draw a 'dot-and-cross' diagram to show the bonding in barium oxide.
			Show outer electrons only.
			[2]
		(iii)	Calculate the number of barium ions in 1.50 g of barium oxide.
			Give your answer in standard form and to three significant figures.
			number of barium ions =[2]
	(b)	Bar	ium chloride, $BaCl_2$, is soluble in water.
		(i)	Compare the electrical conductivities of solid and aqueous barium chloride.
			Explain your answer in terms of the particles involved.


	(11)	Describe the use of aqueous bandin chloride in qualitative analysis.
		[2]
	/:::\	
	(iii)	Hydrated barium chloride can be crystallised from solution.
		Hydrated barium chloride has the formula $BaCl_2 \cdot xH_2O$ and a molar mass of 244.3 g mol ⁻¹ .
		Determine the value of \mathbf{x} in the formula of $\mathrm{BaC}l_2 \cdot \mathbf{x}H_2\mathrm{O}$.
		Show your working.
		x =[2]
(c)	Bar	ium nitride is formed when barium is heated with nitrogen.
` '	(i)	Complete the electron configuration of a nitride ion.
	(')	
		1s ² [1]
	(ii)	Solid barium nitride is reacted with water, forming an alkaline solution A and an alkaline gas B .
		Identify A and B .
		Write an equation, including state symbols, for the reaction.
		A
		В
		[4]

2* A student was asked to carry out an experiment to determine the initial rate of reaction of zinc and hydrochloric acid.

The student plans to collect a total of about $72\,\mathrm{cm}^3$ of hydrogen at RTP and to use an excess of zinc.

The student selects the following apparatus:

- the apparatus shown in the diagram
- 100 cm³ measuring cylinder
- · stop clock
- 2 decimal place balance



outline now the student could carry out the experiment and explain now the results could be processed graphically. Show all working in your calculations.
[6]

3	Alkanes are saturated	hydrocarbons	with the general	formula $C_n H_{2n+2}$.
---	-----------------------	--------------	------------------	--------------------------

(ii)

(a)	A student carries out an experiment to measure the enthalpy change of combustion, A	$\Delta_{c}H$	of
	hexane.		

The student finds that combustion of 1.29 g of hexane changes the temperature of 200 g of water from $20.5\,^{\circ}\text{C}$ to $65.5\,^{\circ}\text{C}$.

(i) Calculate the enthalpy change of combustion, $\Delta_{\rm c}H$, of hexane, in kJ mol⁻¹.

Give your final answer to an **appropriate** number of significant figures.

Δ	_c HkJ mol ⁻¹ [4]
The calculated value of $\Delta_{\rm c}H$ for hexane from book value.	n this experiment is different from the data
Suggest two reasons for this difference.	
1	
2	

© OCR 2017 Turn over


(b) Data book values for the standard enthalpy changes of combustion, $\Delta_{\rm c}H^{\rm e}$, of the first four alkanes are shown in the table.

Alkane	methane	ethane	propane	butane
$\Delta_{\rm c}H^{\rm e}/{\rm kJmol^{-1}}$	-890	-1560	-2219	-2877

(i) The values for the first three alkanes are plotted on the graph below.

Plot the value for butane on the graph.





[1]

(ii)	Use the graph to estimate the energy released during complete combustion of 1.80 g c)f
	pentane.	

Show relevant working below and on the graph.

(c) The equation for the complete combustion of cyclohexane is shown below.

$$\mathsf{C_6H_{12}(I)} + 9\mathsf{O_2(g)} \, \longrightarrow \, \mathsf{6CO_2(g)} + \mathsf{6H_2O(I)}$$

Standard enthalpy changes of formation, $\Delta_{\rm f} H^{\rm e}$, are shown in the table.

Substance	C ₆ H ₁₂ (I)	CO ₂ (g)	H ₂ O(I)
∆ _f H ^e / kJ mol ^{−1}	-156.3	-393.5	-285.8

Calculate the standard enthalpy change of combustion, $\Delta_{\rm c} H^{\rm e},$ of cyclohexane.

$$\Delta_{\rm c} H^{\rm e} =$$
 kJ mol⁻¹ [3]

© OCR 2017 Turn over

- 4 The hydroxyl group, –OH, is responsible for many properties of alcohols.
 - (a) Methanol, CH₃OH, is soluble in water because it has polar bonds.

Pauling electronegativity values for carbon, oxygen and hydrogen are shown below.

Element	Electronegativity
Carbon	2.5
Oxygen	3.5
Hydrogen	2.1

Use a labelled diagram to explain why methanol is soluble in water.

- Use displayed formulae showing one molecule of methanol and one molecule of water.
- Add partial charges δ + and δ to show the **two** most polar bonds in a methanol molecule and the polar bonds in a water molecule.
- Show all lone pairs.
- Label the most important intermolecular bond between the molecules.

(b) Alcohol C is analysed using mass spectrometry.

alcohol C

(i)	Give the systematic name of alcohol C .
	[1]

(ii) The mass spectrum of alcohol **C** is shown below.

Item removed due to third party copyright restrictions.

Write structural formulae for the ions responsible for peak \boldsymbol{X} and peak \boldsymbol{Y} .

X =

Y =

(c)*	Describe the oxidation reactions of butan-1-ol forming an aldehyde and a carboxylic acid.							
	Explain, using a diagram, how the aldehyde can be produced in the laboratory by controlling the reaction conditions.							
	[6]							

5	Pro	panoic acid, CH ₃ CH ₂ COOH, is a member of the homologous series of carboxylic acids.
	(a)	Suggest the general formula for a carboxylic acid.
		[1]

(b) The displayed formula for propanoic acid is shown below.

(i)	State the shape and bond angle around a carbon atom in the alkyl group of propanoic acid. Explain the shape.
	Shape
	Bond angle
	Explanation
	[2]
(ii)	Suggest a value for the C-O-H bond angle in propanoic acid.

[1]


(c)	Compound	D	is	а	neutral	compound	which	is	а	structural	isomer	of	propanoic	acid
	CH ₃ CH ₂ COOl		۱.											

The infrared spectrum of compound $\boldsymbol{\mathsf{D}}$ is shown below.

Item removed due to third party copyright restrictions.
Suggest two possible structures of compound D .
Explain all your reasoning.
F.41

(d)	2-Chloropropanoic acid, CH ₃ CHC1COOH, can be made by reacting propanoic acid wit chlorine in a radical substitution reaction.							
	State the conditions for the reaction.							
		[1]						
	(ii)	Write the overall equation for the reaction.						
		[1]						
(iii) The first step in the reaction mechanism involves homolytic fission of a chlor to form two chlorine radicals.								
		Why is this step an example of homolytic fission?						
		[1]						
((iv)	Write two equations to show the propagation steps in the mechanism for this reaction.						
		Use dots,•, to show the unpaired electrons on radicals.						
		[2]						
	(v)	Draw the displayed formula of the radical formed in the first propagation step.						
		Use a dot,•, to show the position of the unpaired electron.						
		F41						
		[1]						
((vi)	Further substitution forms a mixture of organic products.						
		Draw the structure of an organic product formed from 2-chloropropanoic acid by further substitution.						

- **6** Organic compounds can be prepared in the laboratory using synthetic routes with two or more stages.
 - (a) A student devises a two-stage synthesis of cyclohexene from bromocyclohexane.



(i) Suggest the structure of intermediate E and the reagent(s) and conditions for step 2.

reagent(s) and conditions.....[2]

(ii) The student carries out this synthesis and obtains 1.23 g of pure cyclohexene from 5.50 g of bromocyclohexane.

Calculate the percentage yield of cyclohexene.

Give your final answer to an appropriate number of significant figures.

(b) Cyclohexene is reacted with bromine to prepare the organic compound F.Give the structure of compound F and outline the mechanism for this reaction.Include curly arrows, charges and relevant dipoles.

[4]

END OF QUESTION PAPER

ADDITIONAL ANSWER SPACE

If additional must be cle	al space is required, you should use the following lined page(s). The early shown in the margin(s).	The question number(s)
	D	



Copyright Information

OCR is committed to seeking permission to reproduce all third-party content that it uses in its assessment materials. OCR has attempted to identify and contact all copyright holders whose work is used in this paper. To avoid the issue of disclosure of answer-related information to candidates, all copyright acknowledgements are reproduced in the OCR Copyright Acknowledgements Booklet. This is produced for each series of examinations and is freely available to download from our public website (www.ocr.org.uk) after the live examination series.

If OCR has unwittingly failed to correctly acknowledge or clear any third-party content in this assessment material, OCR will be happy to correct its mistake at the earliest possible opportunity.

For queries or further information please contact the Copyright Team, First Floor, 9 Hills Road, Cambridge CB2 1GE.

OCR is part of the Cambridge Assessment Group; Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.