

Mark Scheme (Results)

Summer 2022

Pearson Edexcel International GCSE In Physics (4PH1) Paper 1P

## **Edexcel and BTEC Qualifications**

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at <a href="https://www.edexcel.com">www.edexcel.com</a> or <a href="https://www.edexcel.com">www.edexcel.com</a>, you can get in touch with us using the details on our contact us page at <a href="https://www.edexcel.com/contactus">www.edexcel.com/contactus</a>.

## Pearson: helping people progress, everywhere

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: <a href="https://www.pearson.com/uk">www.pearson.com/uk</a>

Summer 2022
Question Paper Log Number P70951A
Publications Code 4PH1\_1P\_2206\_MS
All the material in this publication is copyright
© Pearson Education Ltd 2022

## **General Marking Guidance**

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded.
   Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

Question number	Answer	Notes	Marks
1 (a)		comet	
(i)	centre of S marked in the grey region shown;	allow if centre of S marked in a similar position on the other side of the drawn orbit	1
(ii)	closed path drawn with curve; total size approximately twice size of original diagram;	curve should extend at least as far as the start of the word "diagram" in sentence below i.e.	2
(iii)	X marked on part of path closest to candidate's S;	reject if another contradictory cross drawn	1
(b)	<ul> <li>any three from:</li> <li>MP1. comet's orbit is more elliptical/oval;</li> <li>MP2. (orbital) path of comet is longer;</li> <li>MP3. idea that speed of comet varies but speed of planet does not;</li> <li>MP4. both planet and comet orbit a star;</li> <li>MP5. idea that orbital radius of comet varies but orbital radius of planet does not;</li> <li>MP6. plane of comet's orbit different to plane of planet's orbit;</li> <li>MP7. star is at centre of planet orbit but not at centre of comet orbit;</li> </ul>	comparison needs to be seen allow planet's orbit is more circular allow distance for path allow period of orbit of comet is longer  allow the Sun for star  allow (star's) gravitational force is constant for planet but varies for comet	3

Total for Question 1 = 7 marks

Question number	Answer	Notes	Marks
2 (a)	C (number of protons in the nucleus);  A is incorrect because electrons are not found in the B is incorrect because this is deduced from the mass ID is incorrect because this is the mass number		1
(b)	D (number of protons and neutrons in the nucleus);  A is incorrect because electrons are not found in the B is incorrect because this is determined from the manumber itself C is incorrect because this is the atomic number		1
(c)	D (atoms with the same number of protons but a difference of a difference of protons but a difference of a difference of protons but a difference of a differe	umber of protons per of neutrons to each other	1
(d)	A (adding an electron);  B is incorrect because this will change the element ar C is incorrect because this will create a positively change incorrect because this will change the element		1
(e)	B (radioactive decay happens at random);  A is incorrect because this is a consequence of radioa C is incorrect because it does not explain the random D is incorrect because this explains why radioactive d	nature of radioactive decay	1
(f)	A (becquerel (Bq);  B is incorrect because this is the unit for charge C is incorrect because this is the unit for energy D is incorrect because this is the unit for power		1

Question number	Answer	Notes	Marks
3 (a)	40 000 {waves / vibrations / oscillations}; in 1 second / unit time;	allow 40 000 wavelengths / wavefronts allow per second	2
(b) (i)	(wave) speed = frequency × wavelength;	allow standard symbols and rearrangements e.g. $\lambda = v / f$ condone s for speed ignore incorrect symbols e.g wl for wavelength, ws for wave speed	1
(ii)	substitution; rearrangement; evaluation; e.g. $345 = 40 (000) \times \lambda$ $\lambda = 345/40 (000)$ $(\lambda =) 8.6 \times 10^{-3} (m)$	POT error = -1 correct answer with no working gains full marks 8.6, 8.62, 8.63, 8.625 = 2 marks allow $9 \times 10^{-3}$ , $8.62 \times 10^{-3}$ , $8.63 \times 10^{-3}$	3
(iii)	<pre>use of speed = distance/time seen or implied;  factor of 2 seen; substitution;</pre>	10 <sup>-3</sup> , 8.625 × 10 <sup>-3</sup> (m)  allow any rearrangement or in correct symbols allow d, s, x for distance allow v for speed condone s for speed	4
	e.g. speed = distance/time distance should be doubled 345 = 4.70 ÷ time (time =) 0.0136 (s)	0.007, 0.0068 scores 3 marks (no factor of 2 used)  allow 0.01, 0.014, 0.01362 (s)	
(c)	idea that the person can see/know what (distance) they are measuring/eq;	e.g. marks the position that (distance) is being measured to	1

(d)	any three from:	allow points shown on a labelled	3
(d)	any three from:	allow points shown on a labelled	3
		diagram	
		allow higher level ideas i.e. light	
		can be polarised but sound can't	
		allow reverse argument (RA) for	
		all marking points	
	MP1. sound is longitudinal light is	can be given from diagram	
	transverse;	showing longitudinal and	
		transverse waves	
	MP2, sound vibrations are in the direction	allow oscillations, displacements	
	of travel, light vibrations are	for vibrations	
	perpendicular to the direction of	allow direction of energy	
	travel;	transfer for direction of travel	
	MP3. light is electromagnetic, sound is not;	allow sound is mechanical, light	
	Mi 5. tight is electromagnetic, sound is not,	is not	
	MP4. light can travel through a	allow sound needs a medium but	
	1		
	vacuum/space, sound cannot;	light does not	
	MP5. light travels (much) faster (in air)	allow quoted speeds for	
	than sound;	comparison	
	MP6. light slows down in denser medium		
	but sound speeds up;		

Total for Question 3 = 14 marks

Answer	Notes	Marks
suitable linear scale chosen (>50% of grid used);  axes labelled with quantities and unit;  all bar plotting correct to nearest half square;	ignore orientation of axes do not accept multiples of 3 for scale condone missing "planets" label if individual planets are written with bars reject if non-linear scale used  bars do not need to be separated from each other or in the same order as the table  condone stick graphs	3
plan	nets	
any one from: different masses; different sizes; different densities;	ignore different orbital radius ignore weight allow different volume, radii, diameter ignore different (surface) area	1
	suitable linear scale chosen (>50% of grid used);  axes labelled with quantities and unit;  all bar plotting correct to nearest half square;  25  20  3.7  Mercury Verus Turking different masses; different sizes;	suitable linear scale chosen (>50% of grid used);  axes labelled with quantities and unit;  all bar plotting correct to nearest half square;  bars do not need to be separated from each other or in the same order as the table  condone stick graphs  25  20  8.7  12.7  Mercury Verus Turkin Satur Vronus Negroes  any one from: different masses; different sizes;  ignore different orbital radius ignore weight allow different volume, radii, diameter ignore different (surface) area

(c)	as orbital radius increases, the orbital speed decreases;	condone idea that they are inversely proportional ignore negative correlation	1
(d)	substitution into rearrangement; evaluation; $v = \frac{2 \times \pi \times r}{T}$ ; evaluation;	-1 for POT error allow 1s.f. answer if supported by working	3
	e.g. $47.4 = 2 \times \pi \times 57.9 \times 10^6 / T$ $T = 2 \times \pi \times 57.9 \times 10^6 / 47.4$ $(T =) 7.68 \times 10^6$ (s)	allow 7.7×10 <sup>6</sup> , 7.67×10 <sup>6</sup> , 7675 030.154 (s)	

Total for Question 4 = 8 marks

Question number	Answer	Notes	Marks
5 (a)	any suitable natural source; e.g. (the) Sun, cosmic rays, rocks, (named) food, radon etc.	reject if contradicted by a list allow named radioactive isotopes e.g. carbon-14, uranium-235, uranium-238 ignore "space", cosmic microwave background radiation (CMBR)	1
(b) (i	<ul> <li>MP1. alpha has more mass;</li> <li>MP2. alpha has more charge;</li> <li>MP3. alpha is positive and beta is negative;</li> <li>MP4. alpha has shorter range (in air);</li> <li>MP5. alpha is slower;</li> <li>MP6. alpha is less penetrating;</li> <li>MP7. alpha is more ionising;</li> <li>MP8. alpha is a helium nucleus but beta is an electron;</li> <li>evidence of trying to balance nuclear equation; correct number of alphas; correct number of betas;</li> <li>e.g.</li> <li>90 = 86 + 2α</li> </ul>	allow RA throughout allow alpha is heavier  allow oppositely charged  allow alpha stopped by air/paper and beta stopped by aluminium/thin metal  allow alpha is 2 protons and 2 neutrons but beta is an electron  e.g. 232 - 220 = 12 also gains first mark also gains first mark  this balances atomic number despite mass number not balancing (if no beta was present)	3
(c)	<pre>number of alpha = 3 number of beta = 2  any three from: MP1. (alpha) can cause cell mutation / cancer; MP2. idea that alpha is only dangerous when inside body; MP3. alpha is blocked by skin / few cm of air; MP4. thorium can only cause irradiation (since it remains in work surface); MP5. radon / gas can cause (both) contamination (and irradiation) (since it can go inside body / food); MP6. radon / gas can be inhaled / enter body; MP7. thorium cannot enter body;</pre>	allow both (thorium and radon) can cause irradiation	3

Question number	Answer	Notes	Marks
6 (a)	(quantity that has both) magnitude and direction;	allow size, value, number or amount for magnitude ignore scale for magnitude	1
(b)	vertically downward arrow drawn; labelled weight;  arrow drawn in opposite direction to velocity; labelled air resistance / drag / air friction;	ignore starting points and lengths of arrows 3 marks max. if more than two arrows drawn judge by eye dependent on previous mark being awarded (DOP) allow gravitational force, force of gravity ignore gravity, gravitational field strength, gfs  DOP but allow if arrow pointing to the left condone wind resistance	4

Total for Question 6 = 5 marks

	Questi numb		Answer	Notes	Marks
7	(a)		(current is the) rate of flow of charge;	allow amount of charge passing per unit time/second ignore rate of flow of electrons	1
	(b)	(i)	idea that LED shows that current is present / fuse is still working;	allow idea that it shows when the fuse has blown	1
		(ii)	correct symbol for ammeter placed in series with fuse;		1
	(c)	(i)	0.1 (A);		1
		(ii)	0.6 (A);		1
		(iii)	<ul> <li>lamp 3;</li> <li>any two from:</li> <li>(because) power = current × voltage;</li> <li>voltage is the same for all lamps;</li> <li>current is greatest;</li> </ul>	allow ECF from (c)(i) if I <sub>1</sub> given as greater than 0.3(A) leading to lamp 2 being brightest allow P = V <sup>2</sup> / R allow resistance is lowest	3

Total for Question 7 = 8 marks

Question	Answer	Notes	Marks
number 8 (a)	idea that energy can't be created or	both created and destroyed / eq	1 1
	destroyed;	need to be seen ignore lost, removed for destroyed	
(b) (i)	efficiency = $\frac{\text{useful (energy) output}}{\text{total (energy) output}}$ ;	ignore ×100(%) allow rearrangements allow total (energy) input for total (energy) output	1
(ii)	substitution; rearrangement; evaluation of useful energy output; subtraction from total to find wasted energy output;  e.g. 0.15 = E <sub>useful</sub> / 165 E <sub>useful</sub> = 0.15 × 165	allow 24.75 (J) allow 140.25 (J)  allow alternative method of determining 85% of energy wasted, leading to correct final answer e.g.  % wasted = 100 - 15 = 85 0.85 = E <sub>wasted</sub> / 165	4
	(E <sub>useful</sub> =) 25 (J) (E <sub>wasted</sub> =) 165 - 25 = 140 (J)	(E <sub>wasted</sub> =) 0.85 × 165 (E <sub>wasted</sub> =) 140 (J)	
(iii)	diagram to show standard single input with two output arrows; input and output arrows labelled correctly; scale approximately correct;	arrows can be in any orientation e.g. both useful and wasted arrows drawn horizontally allow elastic, input, total for initial arrow kinetic, useful (output) for narrower output thermal/heat, wasted (output) for wider output but apply ECF from (b)(ii) allow ECF from (b)(ii) judge by eye - wasted arrow should be at least 4× wider than useful arrow and they should sum to approximately the width of the input	3
	e.g.  elastic  thermal		

(c) (i)	45 (J);		1
(ii)	substitution into work done = force × distance; rearrangement; evaluation;  e.g. 45 = force × 7.5 force = 45 / 7.5		3
	(force =) 6.0 (N)	allow 6 (N)	

Total for Question 8 = 13 marks

Question	Answer	Notes	Marks
9 (a) (i)	pressure (difference) = height × density × g;	allow standard symbols and rearrangements e.g. $h = p / (\rho \times g)$ condone d for density	1
(ii)	substitution; rearrangement; evaluation in cm;  e.g.	allow g = 9.81, 9.8  -1 for POT error but check carefully due to value of g 0.23 (cm) gains 2 marks	3
	2300 = h × 1000 × 10 h = 2300 / (1000 × 10) (h =) 23 (cm)	allow 23.46, 23.44 (cm)	
(b) (i)	line drawn of similar curved path and aiming towards ground to the left of path from hole B;	reject if curve drawn upwards at any point	1
(ii)	any two from:  MP1. pressure is lower at A; MP2. force acting on water at A is less; MP3. water leaves the bottle at a lower speed/KE at A;  OR  any two from: MP1. initial velocity of water is horizontal;	allow "near top of bottle"/eq for A allow RA allow RA allow RA	2
	MP2. (force of) gravity acts on the water; MP3. water accelerates downwards;	horizontally	
(iii)	any three from:  MP1. idea that path from C is identical / symmetrical to path from B;  MP2. (because) pressure is the same;  MP3. (initial) speed of water is the same;  MP4. idea that pressure acts (equally) in all directions;	ignore holes at same height above ground allow path is the same (as B) allow force acting on water is the same allow (initial) KE of water is the same	3
(iv)	any one from:  MP1. idea that it allows air to enter the bottle (as water level falls);  MP2. idea that it maintains equal pressure between air inside and air outside the bottle;  MP3. idea that no water would come out of the holes otherwise;	Total for Question 9 - 11 mark	1

Question number	Answer	Notes	Marks	
10 (a)	substitution into v <sup>2</sup> = u <sup>2</sup> + 2as; rearrangement; evaluation;	allow alternative method involving changing GPE lost = KE gained e.g. sub into mgh = ½mv²  allow use of 9.8, 9.81 for g	3	
	e.g. $v^2 = 0 + (2 \times 10 \times 18)$ $v = \sqrt{360}$			
	(v =) 19 (m/s)	allow 18.8, 18.78, 18.79,18.97 (m/s)		
(b) (i)	kinetic energy = $\frac{1}{2} \times \text{mass} \times \text{speed}^2$ ;	allow rearrangements and standard symbols e.g. KE = ½ × m × v <sup>2</sup>	1	
(ii)	substitution; evaluation; e.g.	allow ECF from (a) allow alternative method involving changing GPE lost = KE gain allow 370 000-372 000 from g = 9.8, 9.81	2	
	(KE =) 0.5 × 2100 × 19 <sup>2</sup> (KE =) 380 000 (J)	allow 400 000, 378 000, 379 000 1 mark only for not converting tonnes to kg e.g. KE = 380 (J), 378 (J)		
(c)	idea that energy is transferred from a gravitational (store) to a kinetic (store);	allow answers in terms of types of energy rather than stores and transfers allow GPE to KE	4	
	idea that energy is transferred to a thermal (store) of {car / shaft / surroundings};	allow heat for thermal		
	idea that energy is transferred mechanically;	allow energy transferred due to a named force e.g. gravity, friction		
	idea that energy is transferred by radiation / by heating;	allow energy transferred as sound		

Question number	Answer	Notes	Marks
11 (a)	any two from: MP1. height is <u>independent variable</u> ; MP2. at least five different heights tested; MP3. range is <u>dependent variable</u> ;		6
	with any two from: MP4. one control variable named; MP5. second control variable named; MP6. idea of repeating process at each height to find mean;	e.g. launch speed/launch force, angle of launch, same ball allow repeating process at each height to identify anomalies	
	AND: MP7. ruler used to measure height / range; MP8. suitable method to see where ball lands;	allow tape measure condone metre stick e.g. record video (and playback in slow motion) cover ball in paint, material on floor to show landing point etc.	
(b) (i)	smooth curve within one small square of data points;	ignore extrapolation of curve beyond the points take care the curve is not dot to dot straight lines	1
(ii)	height reading consistent with curve of best fit;	allow 0.15-0.25 (m) allow ECF from (b)(i)	1
(iii)	idea that (both) variables are continuous;	allow results/data are continuous	1
(iv)	pair of readings taken from graph; substitution into given formula; rearrangement; evaluation to at least 1 decimal place;	allow data points or readings taken from candidate's curve allow ECF from (b)(i) and (b)(ii) 3 marks max. for reverse	4
	e.g. range = 1.92m when height = 2.0m 1.92 = launch speed $\times \sqrt{\frac{2.0}{5}}$	argument e.g. using speed of 3 m/s to calculate height or range	
	launch speed = 1.92 / $\sqrt{0.4}$ (launch speed =) 3.0 (m/s)	allow range 2.9-3.1 (m/s) unless ECF from (b)(i)	

Question number			Answer	Notes	Marks
12	(a)		2.8 (mA);	allow -2.8 (mA)	1
	(b)	(i)	same value as (a) but opposite polarity e.g2.8 (mA);	allow ecf from (a)	1
		(ii)	reading larger than value given in (a);	ignore polarity	1
	(c)		any one from: MP1. longer needle; MP2. more turns in coil; MP3. stronger magnet; MP4. weaker return spring;	allow idea of moving scale further up the needle ignore bigger coil, more coils ignore bigger magnet	1
	(d)	(i)	A (accuracy);  B is incorrect because no repeats are being taken C is incorrect because no repeats are being taken D is incorrect because the scale of the ammeter is the same		1
		(ii)	idea of subtracting/adding the difference/error from the measurement;	allow subtract/add a quoted value of current even if value is incorrect	1

Total for Question 12 = 6 marks