For more awesome GCSE and A level resources, visit us at <a href="https://www.savemyexams.co.uk/">www.savemyexams.co.uk/</a>

## **Vector Geometry**

## **Question Paper**

Level	Pre U
Subject	Maths
Exam Board	Cambridge International Examinations
Topic	Vector Geometry
Booklet	Question Paper

Time Allowed: 52 minutes

Score: /43

Percentage: /100

**Grade Boundaries:** 

## Save My Exams! - The Home of Revision

For more awesome GCSE and A level resources, visit us at www.savemyexams.co.uk/

The points A and B have position vectors **a** and **b** relative to an origin O, where  $\mathbf{a} = 5\mathbf{i} + 4\mathbf{j} - 2\mathbf{k}$  and  $\mathbf{b} = -7\mathbf{i} + 3\mathbf{j} + \mathbf{k}$ .

- (i) Find the length of AB. [3]
- (ii) Use a scalar product to find angle *OAB*. [4]
- 2 Two intersecting straight lines have equations

$$\frac{x-5}{4} = \frac{y-11}{3} = \frac{z-7}{-5}$$
 and  $\frac{x-9}{-2} = \frac{y-4}{1} = \frac{z+4}{4}$ .

[6]

Find the coordinates of their point of intersection.

3 Vectors **u** and **v** are given by  $\mathbf{u} = \begin{pmatrix} 4 \\ 6 \end{pmatrix}$  and  $\mathbf{v} = \begin{pmatrix} -3 \\ 2 \end{pmatrix}$ .

(i) Find 
$$\mathbf{u} + \mathbf{v}$$
 and  $\mathbf{u} - \mathbf{v}$ . [2]

(ii) Show that 
$$|\mathbf{u} + \mathbf{v}| = |\mathbf{u} - \mathbf{v}|$$
. [2]

- 4 Three points A, B and C have coordinates (1, 0, 7), (13, 9, 1) and (2, -1, -7) respectively.
  - (i) Use a scalar product to find angle *ACB*. [5]
  - (ii) Hence find the area of triangle ACB. [2]
  - (iii) Show that a vector equation of the line *AB* is given by  $\mathbf{r} = \mathbf{i} + 7\mathbf{k} + \lambda(4\mathbf{i} + 3\mathbf{j} 2\mathbf{k})$ , where  $\lambda$  is a scalar parameter.
- 5 (i) Given that the point (-1, -2, 4) lies on both the lines

$$\mathbf{r} = \begin{pmatrix} 2 \\ -3 \\ a \end{pmatrix} + \lambda \begin{pmatrix} -3 \\ 1 \\ 1 \end{pmatrix}$$
 and  $\mathbf{r} = \begin{pmatrix} 2 \\ 4 \\ b \end{pmatrix} + \mu \begin{pmatrix} -1 \\ -2 \\ 1 \end{pmatrix}$ ,

find a and b. [3]

(ii) Find the acute angle between the lines. [4]

## Save My Exams! - The Home of Revision

For more awesome GCSE and A level resources, visit us at www.savemyexams.co.uk/

- The points A and B have position vectors  $\mathbf{i} \mathbf{j} + \mathbf{k}$  and  $2\mathbf{i} + \mathbf{j} + 3\mathbf{k}$  respectively, relative to the origin O. The point C is on the line OA extended so that  $\overrightarrow{AC} = 2\overrightarrow{OA}$  and the point D is on the line OB extended so that  $\overrightarrow{BD} = 3\overrightarrow{OB}$ . The point X is such that OCXD is a parallelogram.
  - (i) Show that a vector equation of the line AX is  $\mathbf{r} = \mathbf{i} \mathbf{j} + \mathbf{k} + \lambda(5\mathbf{i} + \mathbf{j} + 7\mathbf{k})$  and find an equation of the line CD in a similar form. [5]
  - (ii) Prove that the lines AX and CD intersect and find the position vector of their point of intersection. [4]